

VETSCRIPTION SURE SHOT 2X - pyrantel pamoate liquid
Sergeant's Pet Care Products, Inc.

INDICATIONS FOR USE

Sure Shot 2X[®] suspension is highly palatable formulation intended as a single treatment for the removal of large rounds (*Toxocara canis* and *Toxocaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. Sure Shot 2X[®] suspension may also be used to prevent re-infestation of *Toxocara canis* in puppies and adult dogs and lactating bitches after whelping. The presence of these parasites should be confirmed by laboratory fecal examination. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

DOSAGE AND ADMINISTRATION

IMPORTANT: SHAKE WELL BEFORE USING

For maximum control and prevention of re-infestation, it recommended that puppies be treated 2, 3, 4, 6, 8 and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* re-infestation.

Administer one full teaspoon (5 mL) for each 10 pounds of body weight (2.27 mg base per lb. of body weight). Although most dogs have been observed to find this formulation very palatable and willingly consume it undiluted, it may be necessary to mix a small quantity of formula in the dog's normal ration of dog food to encourage consumption. Fasting prior to and after treatment is not necessary.

CAUTION

This product is a suspension and as such will separate. To ensure uniform resuspension and to achieve proper dosage, it is extremely important that the product be shaken thoroughly before every use.

KEEP OUT OF REACH OF CHILDREN

RECOMMENDED STORAGE

Store below 30°C (86°F)

ACTIVE INGREDIENT

4.54 mg of pyrantel base as pyrantel pamoate per mL

HOW SUPPLIED

Sure Shot 2X[®] - Net Contents: 2 fl. oz. (60 mL) NDC 21091-408-02

RECOMMENDATION

Dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2 to 4 weeks after first treatment.

PACKAGING COMPONENTS

Sergeant's Vetscription[®] Sure Shot 2X[®] Liquid Wormer (pyrantel pamoate)
FOR ANIMAL USE ONLY

Read Accompanying Professional Information

ANADA 200-007, Approved by FDA

Distributed by:

Sergeant's Pet Care Products, Inc., Omaha NE 68130

www.sergeants.com

INDICATIONS FOR USE:

To prevent reinfestation of *Toxocara canis* in puppies and adult dogs and lactating bitches after whelping.

For the removal of large roundworms, (*Toxocara canis* and *Toxascaris leonina*) and hookworms, (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies.

The presence of these parasites should be confirmed by laboratory fecal examination. Consult your veterinarian for assistance in the diagnosis treatment and control of parasitism.

WARNING: KEEP OUT OF THE REACH OF CHILDREN.

Read Accompanying Professional Information

Recommended Storage: STORE BELOW 30°C (86°F)

Distributed by: Sergeant's Pet Care Products, Inc., Omaha, NE 68130

Visit us at www.sergeants.com

P006005-2

RRAV

**DO NOT PRINT
UNCOATED AREA**

for puppies and dogs
up to 120 lbs.


Sergeant's
VeTSCRIPTION

Sure Shot '2X'


(pyrantel pamoate)

Liquid Wormer

Removes

 **hookworms**

(*Ancylostoma caninum*, *Uncinaria stenocephala*)

 **large roundworms**

(*Ascarids*)

Active ingredient 4.54 mg of pyrantel

base as pyrantel pamoate per mL

FOR ANIMAL USE ONLY

See back panel for complete use directions



Net Contents: 2 fl oz (60 mL)
NADA 200-007 Approved by FDA

IMPORTANT: SHAKE WELL BEFORE USING DIRECTIONS FOR USE:

For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T.canis* reinfestation. Administer one full teaspoonful (5 mL) for each 10 lb of body weight.

For the removal of large roundworms (ascarids) and hookworms. Administer one full teaspoonful (5 mL) for each 10 lbs. of body weight. It is not necessary to withhold food prior to treatment.

If medication is to be dispensed, client can be advised that dogs usually find this wormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption.

The presence of large roundworms (ascarids) and hookworms should be confirmed by a lab fecal examination.

It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2 to 4 weeks after first treatment.

